arXiv:1612.08819v1 [math.FA] 28 Dec 2016

# CHARACTERIZATIONS OF THE BMO AND LIPSCHITZ SPACES VIA COMMUTATORS ON WEAK LEBESGUE AND MORREY SPACES

DINGHUAI WANG, JIANG ZHOU* AND WENYI CHEN

Abstract. We prove that the weak Morrey space $WM^p_q$ is contained in the Morrey space $M^p_{q_1}$ for $1 \le q_1 < q \le p < \infty$. As applications, we show that if the commutator $[b, T]$ is bounded from $L^p$ to $L^{p,\infty}$ for some $p \in (1, \infty)$, then $b \in$ BMO, where $T$ is a Calderón-Zygmund operator. Also, for $1 < p \le q < \infty$, $b \in$ BMO if and only if $[b, T]$ is bounded from $M^p_q$ to $WM^p_q$. For $b$ belonging to Lipschitz class, we obtain similar results.

## 1. Introduction

Let $T$ be a Calderón-Zygmund operator defined by

$$Tf(x) = \text{p.v.} \int_{\mathbb{R}^n} K(x-y)f(y)dy,$$

where the kernel $K(x) = \frac{\Omega(x)}{|x|^n}$ satisfies the following conditions:

(i) $\Omega$ is homogeneous of degree zero on $\mathbb{R}^n$, i.e., $\Omega(\lambda x) = \Omega(x)$ for all $\lambda > 0$ and $x \in \mathbb{R}^n$;

(ii) $\Omega \in C^\infty(\mathbb{S}^{n-1})$ and $\int_{\mathbb{S}^{n-1}} \Omega(x)dx = 0$.

A locally integrable function $b$ belongs to the BMO space if $b$ satisfies

$$\|b\|_* := \sup_Q \frac{1}{|Q|} \int_Q |b(x) - b_Q|dx < \infty,$$

where $b_Q := \frac{1}{|Q|} \int_Q b(x)dx$ and the supremum is taken over all cubes $Q$ in $\mathbb{R}^n$. A well known result of Coifman, Rochberg and Weiss [2] states that the commutator

$$[b, T](f) := bT(f) - T(bf)$$

is bounded on some $L^p$, $1 < p < \infty$, if and only if $b \in$ BMO. An interesting question is raised. Is $b$ in BMO if $[b, T]$ is of weak type $(p, p)$ for some $p \in (1, \infty)$? We will give an affirmative answer in this paper.

*2000 Mathematics Subject Classification.* Primary 47B47, 46B04.

*Key words and phrases.* BMO space, Characterization, Commutator, Lipschitz space, Morrey space.

The research was supported by National Natural Science Foundation of China (Grant No.11261055).

* Corresponding author, Email: zhoujiangshuxue@126.com.

For $1 \le q \le p < \infty$, we say that a function $f$ belongs to Morrey space $M^p_q$ if

$$\|f\|_{M^p_q} := \sup_{x\in\mathbb{R}^n, r>0} \frac{1}{|Q(x,r)|^{1/q-1/p}} \bigg( \int_{Q(x,r)} |f(y)|^q dy \bigg)^{1/q} < \infty;$$

a function $f$ belongs to weak Morrey space $WM^p_q$ if

$$\|f\|_{WM^p_q} := \sup_{x\in\mathbb{R}^n, r>0} \frac{1}{|Q(x,r)|^{1/q-1/p}} \sup_{\lambda>0} \bigg( \lambda^q |\{y \in Q(x,r) : |f(y)| > \lambda\}| \bigg)^{1/q} < \infty.$$

Morrey spaces describe local regularity more precisely than $L^p$ spaces and can be seen as a complement of $L^p$. In fact, $L^p = M^p_p \subset M^p_q$ and $WM^p_p = L^{p,\infty}$ for $1 \le q \le p < \infty$. In 1997, Ding [4] showed that $b$ is in BMO if and only if the commutator $[b, T]$ of Calderón-Zygmund operator $T$ is bounded on Morrey spaces. We will demonstrate here that $b \in$ BMO if and only if $[b, T]$ is weak bounded on Morrey spaces.

Another subject of this paper is to consider the characterizations of Lipschitz functions via commutators. For $0 < \alpha < 1$, the Lipschitz space $\mathrm{Lip}_\alpha$ is the set of functions $f$ such that

$$\|f\|_{\mathrm{Lip}_\alpha} := \sup_{\substack{x,h\in\mathbb{R}^n \\ h\neq 0}} \frac{|f(x+h) - f(x)|}{|h|^\alpha} < \infty.$$

In 1978, Janson [5] proved that, for $1 < p < q < \infty$ with $1/q = 1/p - \alpha/n$, $b \in \mathrm{Lip}_\alpha$ if and only if $[b, T]$ is bounded from $L^p$ to $L^q$. In 1995, using Sobolev-Besov embedding, Paluszyński [8] obtained that, for $1 < p < \infty$ and $0 < \alpha < 1$, $b \in \mathrm{Lip}_\alpha$ if and only if $[b, T]$ is bounded from $L^p$ to the homogeneous Triebel-Lizorkin spaces $\dot{F}^{\alpha,\infty}_p$. Paluszyński's idea was novel for the study about the boundedness of commutators from $L^p$ to $\dot{F}^{\alpha,\infty}_p$ and shed new light on the characterization of Lipschitz space via commutators. Recently, Shi and Lu [10] showed that, for $1 < q \le p < \infty$, $1 < t \le s < \infty$, $\frac{1}{s} = \frac{1}{p} - \frac{\alpha}{n}$ and $\frac{1}{t} = \frac{1}{q} - \frac{\alpha}{n}$, $b \in \mathrm{Lip}_\alpha$ if and only if $[b, T]$ is bounded from $M^p_q$ to $M^s_t$. We are going to show that, for $1 \le p < q < \infty$ with $1/q = 1/p - \alpha/n$, $b \in \mathrm{Lip}_\alpha$ if and only if $[b, T]$ is bounded from $L^p$ to $L^{q,\infty}$. Also, for $1 \le q \le p < \infty$, $1 < t \le s < \infty$, $\frac{1}{s} = \frac{1}{p} - \frac{\alpha}{n}$ and $\frac{1}{t} = \frac{1}{q} - \frac{\alpha}{n}$, $b \in \mathrm{Lip}_\alpha$ if and only if $[b, T]$ is bounded from $M^p_q$ to $WM^s_t$.

Throughout this paper, the letter $C$ denotes constants which are independent of main variables and may change from one occurrence to another.

## 2. Characterization of BMO space via commutators

In this section, we characterize BMO space via the boundedness of commutator on (weak) Lebesgue spaces or (weak) Morrey spaces. First of all, we compare with Morrey spaces and weak Morrey spaces.

It is clear that $M^p_{q_1}$ is contained in $WM^p_{q_2}$ and $\|\cdot\|_{WM^p_{q_2}} \le C\|\cdot\|_{M^p_{q_1}}$ if $1 \le q_2 \le q_1 \le p < \infty$. However, for $1 \le q_1 < q_2 \le p < \infty$, one has the reverse inequality as follows.

**Theorem 2.1.** *If $1 \le q_1 < q_2 \le p < \infty$, then $WM^p_{q_2} \subset M^p_{q_1}$ and $\|f\|_{M^p_{q_1}} \le 2\big(\frac{q_1}{q_2-q_1}\big)^{\frac{1}{q_2}}\|f\|_{WM^p_{q_2}}$.*

*Proof.* Let $f \in WM^p_{q_2}$. Given a cube $Q \subset \mathbb{R}^n$ and $\lambda > 0$,

$$\frac{1}{|Q|^{1/q_2-1/p}}\big(\lambda^{q_2}|\{x \in Q : |f(x)| > \lambda\}|\big)^{1/q_2} \le \|f\|_{WM^p_{q_2}};$$

that is,

$$|\{x \in Q : |f(x)| > \lambda\}| \le \|f\|^{q_2}_{WM^p_{q_2}}|Q|^{1-q_2/p}\lambda^{-q_2}.$$

Choose

$$N = \|f\|_{WM^p_{q_2}}|Q|^{-1/p}\Big(\frac{q_1}{q_2-q_1}\Big)^{1/q_2}.$$

Thus,

$$\begin{aligned}\int_Q |f(x)|^{q_1}dx &= q_1\int_0^\infty \lambda^{q_1-1}|\{x \in Q : |f(x)| > \lambda\}|d\lambda \\ &\le q_1\int_0^N \lambda^{q_1-1}|Q|d\lambda + q_1\int_N^\infty \lambda^{q_1-1}\|f\|^{q_2}_{WM^p_{q_2}}|Q|^{1-q_2/p}\lambda^{-q_2}d\lambda \\ &= |Q|N^{q_1} + \frac{q_1}{q_2-q_1}\|f\|^{q_2}_{WM^p_{q_2}}|Q|^{1-q_2/p}N^{q_1-q_2},\end{aligned}$$

which gives

$$\Big(\int_Q |f(y)|^{q_1}dy\Big)^{1/q_1} \le 2\|f\|_{WM^p_{q_2}}|Q|^{1/q_1-1/p}\Big(\frac{q_1}{q_2-q_1}\Big)^{1/q}.$$

Then

$$\|f\|_{M^p_{q_1}} \le 2\Big(\frac{q_1}{q_2-q_1}\Big)^{1/q_2}\|f\|_{WM^p_{q_2}}$$

and the lemma follows. □

**Remark 2.1.** For $1 \le q_1 < q_2 < p < \infty$. Kozono and Yamazaki [7, Lemma 1.7] proved that $\|f\|_{M^p_{q_2}} \le C\|f\|_{L^{p,\infty}}$. The above Theorem 2.1 yields immediately

$$\|f\|_{M^p_{q_1}} \le C\|f\|_{WM^p_{q_2}} \le C\|f\|_{M^p_{q_2}} \le C\|f\|_{WM^p_p} = C\|f\|_{L^{p,\infty}}.$$

Now we return to our first subject.

**Theorem 2.2.** *Let $1 < p < \infty$. The following statements are equivalent:*

(1) $b \in \mathrm{BMO}$*;*
(2) $[b,T]$ *is a bounded operator from $L^p$ to $L^p$;*

(3) $[b,T]$ *is a bounded operator from* $L^p$ *to* $L^{p,\infty}$.

**Theorem 2.3.** *Let* $1 < q \le p < \infty$. *The following statements are equivalent:*

(1) $b \in \mathrm{BMO}$;
(2) $[b,T]$ *is a bounded operator from* $M^p_q$ *to* $M^p_q$;
(3) $[b,T]$ *is a bounded operator from* $M^p_q$ *to* $WM^p_q$.

*Proof of Theorem 2.2.* The equivalence of (1) and (2) was proved in [3]. By the inequality $\|f\|_{L^{p,\infty}} \le \|f\|_{L^p}$, it is obvious that (2) implies (3).

To show (3) $\Rightarrow$ (1), we use Paluszyński idea given in [8]. For $z_0 \in \mathbb{R}^n \backslash \{0\}$, let $\delta = \frac{|z_0|}{2\sqrt{n}}$ and $Q(z_0,\delta)$ denote the open cube centered at $z_0$ with side length $2\delta$. Then $K(x)^{-1}$ has an absolutely convergent Fourier series

$$\frac{1}{K(x)} = \sum a_m e^{i\langle v_m, x\rangle}$$

with $\sum |a_m| < \infty$, where the exact form of the vectors $v_m$ is unrelated. Then, we have the expansion

$$\frac{1}{K(x)} = \frac{\delta^{-n}}{K(\delta x)} = \delta^{-n} \sum a_m e^{i\langle v_m, \delta x\rangle} \ for\ |x - \frac{z_0}{\delta}| < \sqrt{n}.$$

Given cubes $Q = Q(x_0, r)$ and $Q' = Q(x_0 - rz_1, r)$, if $x \in Q$ and $y \in Q'$, then

$$\Big|\frac{x-y}{r} - \frac{z_0}{\delta}\Big| \le \Big|\frac{x-x_0}{r}\Big| + \Big|\frac{y-(x_0 - \frac{rz_0}{\delta})}{r}\Big| < \sqrt{n}.$$

Let $s(x) = \overline{\mathrm{sgn}(\int_{Q'}(b(x)-b(y))dy)}$. Then

$$\begin{aligned}
&\frac{1}{|Q|}\int_Q |b(x) - b_{Q'}|dx \\
&= \frac{1}{|Q|}\frac{1}{|Q'|}\int_Q \Big|\int_{Q'}(b(x)-b(y))dy\Big|dx \\
&= \frac{1}{|Q|^2}\int_Q\int_{Q'} s(x)(b(x)-b(y))dydx \\
&= \frac{1}{|Q|^2}\int_Q\int_{Q'} s(x)(b(x)-b(y))\frac{r^n K(x-y)}{K(\frac{x-y}{r})}dydx \\
&= \frac{1}{|Q|}\int_{\mathbb{R}^n}\int_{\mathbb{R}^n}(b(x)-b(y))K(x-y)\sum a_m e^{i\langle v_m, \delta\frac{x-y}{r}\rangle}s(x)\chi_Q(x)\chi_{Q'}(y)dydx \\
&= \frac{1}{|Q|}\sum a_m \int_{\mathbb{R}^n}\int_{\mathbb{R}^n}(b(x)-b(y))K(x-y)e^{i\langle v_m, \frac{\delta}{r}x\rangle}s(x)\chi_Q(x)e^{-i\langle v_m, \frac{\delta}{r}y\rangle}\chi_{Q'}(y)dydx.
\end{aligned}$$

Setting $g_m(y) = e^{-i\langle v_m, \frac{\delta}{r}y\rangle}\chi_{Q'}(y)$ and $h_m(x) = e^{i\langle v_m, \frac{\delta}{r}x\rangle}s(x)\chi_Q(x)$, we have

$$\begin{aligned}\frac{1}{|Q|}\int_Q |b(x) - b_{Q'}|dx &= \frac{1}{|Q|}\sum a_m \int_{\mathbb{R}^n}\int_{\mathbb{R}^n}(b(x)-b(y))K(x-y)g_m(y)h_m(x)dydx \\ &= \frac{1}{|Q|}\sum a_m \int_{\mathbb{R}^n}[b,T](g_m)(x)h_m(x)dx.\end{aligned}$$

Choose $q \in (1,p)$. By Remark 2.1,

$$\begin{aligned}\frac{1}{|Q|}\int_Q |b(x)-b_Q|dx &\le \frac{2}{|Q|}\int_Q |b(x)-b_{Q'}|dx \\ &\le \frac{C}{|Q|^{1/p}}\sum |a_m|\|[b,T](g_m)\|_{M^p_q} \\ &\le \frac{C}{|Q|^{1/p}}\sum |a_m|\|[b,T](g_m)\|_{L^{p,\infty}} \\ &\le \frac{C\|g_m\|_{L^p}}{|Q|^{1/p}}\|[b,T]\|_{L^p\to L^{p,\infty}} \\ &\le C\|[b,T]\|_{L^p\to L^{p,\infty}},\end{aligned} \tag{2.1}$$

which yields $b \in \mathrm{BMO}$ and $\|b\|_* \le C\|[b,T]\|_{L^p\to L^{p,\infty}}$. Hence, the proof of Theorem 2.2 is completed. □

*Proof of Theorem 2.3.* We use a same argument as the proof of Theorem 2.2 except choosing $q_1 \in (1,q)$ and replacing (2.1) by

$$\begin{aligned}\frac{1}{|Q|}\int_Q |b(x)-b_Q|dx &\le \frac{2}{|Q|}\int_Q |b(x)-b_{Q'}|dx \\ &\le \frac{C}{|Q|^{1/p}}\sum |a_m|\|[b,T](g_m)\|_{M^p_{q_1}} \\ &\le \frac{C}{|Q|^{1/p}}\sum |a_m|\|[b,T](g_m)\|_{WM^p_q} \\ &\le \frac{C\|g_m\|_{M^p_q}}{|Q|^{1/p}}\|[b,T]\|_{M^p_q\to WM^p_q} \\ &\le C\|[b,T]\|_{M^p_q\to WM^p_q}.\end{aligned}$$

Then $b \in \mathrm{BMO}$ and $\|b\|_* \le C\|[b,T]\|_{M^p_q\to WM^p_q}$. This completes the proof of Theorem 2.3. □

## 3. Characterization of $\mathrm{Lip}_\alpha$ via commutators

We give a lemma that can be used to prove a characterization of Lipschitz functions.

**Lemma 3.1.** *For $0 < \alpha < 1$ and $1 < q \le \infty$,*

$$\|b\|_{\mathrm{Lip}_\alpha} \approx \sup_Q \frac{1}{|Q|^{1+\alpha/n}} \int_Q |b(x) - b_Q| dx$$
$$\approx \begin{cases} \sup_Q \frac{1}{|Q|^{\alpha/n}} \left( \frac{1}{|Q|} \int_Q |b(x) - b_Q|^q dx \right)^{1/q} & \text{if } 1 < q < \infty \\ \operatorname{ess\,sup}_Q \frac{|b(x) - b_Q|}{|Q|^{\alpha/n}} & \text{if } q = \infty \end{cases},$$

*where the supremum is taken over all cubes $Q \subset \mathbb{R}^n$ and $\approx$ means equivalence.*

*Proof.* The first equivalence can be found in [3, pages 14 and 38], and the second equivalence can be found in [6]. □

The first result of this section is

**Theorem 3.2.** *Let $0 < \alpha < 1$, $1 \le p < \frac{n}{\alpha}$ and $1/q = 1/p - \alpha/n$. The following statements are equivalent:*

(1) *$b \in \mathrm{Lip}_\alpha$;*
(2) *$[b, T]$ is a bounded operator from $L^p$ to $L^{q,\infty}$.*

*Proof.* (1) $\Rightarrow$ (2): Let $b \in \mathrm{Lip}_\alpha$. Then

$$\begin{aligned} \big|[b,T](f)(x)\big| &= \left| \int_{\mathbb{R}^n} (b(x) - b(y)) f(y) K(x-y) dy \right| \\ &\le \|b\|_{\mathrm{Lip}_\alpha} \int_{\mathbb{R}^n} \frac{|f(y)|}{|x-y|^{n-\alpha}} dy \\ &\le \|b\|_{\mathrm{Lip}_\alpha} I_\alpha(|f|)(x), \end{aligned} \tag{3.1}$$

which implies

$$\|[b,T]f\|_{L^{q,\infty}} \le \|b\|_{\mathrm{Lip}_\alpha} \|I_\alpha(|f|)\|_{L^{q,\infty}} \le C \|b\|_{\mathrm{Lip}_\alpha} \|f\|_{L^p}.$$

(2) $\Rightarrow$ (1): We follow the method in the proof of Theorem 2.2 except choosing $q_1 \in (1, q)$ and replacing (2.1) by

$$\begin{aligned} \frac{1}{|Q|} \int_Q |b(x) - b_Q| dx &\le \frac{2}{|Q|} \int_Q |b(x) - b_{Q'}| dx \\ &\le \frac{C}{|Q|^{1/q}} \sum |a_m| \|[b,T](g_m)\|_{M^q_{q_1}} \\ &\le \frac{C}{|Q|^{1/q}} \sum |a_m| \|[b,T](g_m)\|_{L^{q,\infty}} \end{aligned}$$

$$\leq \frac{C\|g_m\|_{L^p}}{|Q|^{1/q}}\|[b,T]\|_{L^p\to L^{q,\infty}}$$
$$\leq C|Q|^{\alpha/n}\|[b,T]\|_{L^p\to L^{q,\infty}}.$$

From Lemma 3.1 we conclude that $b\in \mathrm{Lip}_\alpha$ and $\|b\|_{\mathrm{Lip}_\alpha}\leq C\|[b,T]\|_{L^p\to L^{q,\infty}}$. Hence, the proof of Theorem 3.2 is completed. □

Using Theorem 3.2 and Janson's result [5], we immediately have

**Corollary 3.3.** *Let $0<\alpha<1$, $1<p<\frac{n}{\alpha}$ and $1/q=1/p-\alpha/n$. The following statements are equivalent:*

(1) *$b\in \mathrm{Lip}_\alpha$;*
(2) *$[b,T]$ is a bounded operator from $L^p$ to $L^q$;*
(3) *$[b,T]$ is a bounded operator from $L^p$ to $L^{q,\infty}$.*

We give two remarkable results about the boundedness of fractional integral operator $I_\alpha$ on Morrey spaces, where $I_\alpha$ is defined by

$$I_\alpha f(x)=\int_{\mathbb{R}^n}\frac{f(y)}{|x-y|^{n-\alpha}}dy,\qquad x\in\mathbb{R}^n \text{ and } 0<\alpha<n.$$

**Theorem 3.4.** (Peetre [9]) *Let $0<\alpha<n$, $1<q\leq p<\infty$ and $1<t\leq s<\infty$. If $\frac{1}{s}=\frac{1}{p}-\frac{\alpha}{n}$ and $\frac{1}{t}=\frac{1}{q}-\frac{\alpha}{n}$, then $\|I_\alpha f\|_{M^s_t}\leq C\|f\|_{M^p_q}$.*

**Theorem 3.5.** (Adams [1]) *Let $0<\alpha<n$, $1<q\leq p<\infty$ and $1<l\leq s<\infty$. If $\frac{1}{s}=\frac{1}{p}-\frac{\alpha}{n}$ and $\frac{l}{s}=\frac{q}{p}$, then $\|I_\alpha f\|_{M^s_l}\leq C\|f\|_{M^p_q}$.*

We will use the above two theorems to show other characterizations of Lipschitz functions.

**Theorem 3.6.** *Let $0<\alpha<1$, $1\leq q\leq p<\infty$, $1<t\leq s<\infty$, $\frac{1}{s}=\frac{1}{p}-\frac{\alpha}{n}$ and $\frac{1}{t}=\frac{1}{q}-\frac{\alpha}{n}$. The following statements are equivalent:*

(1) *$b\in \mathrm{Lip}_\alpha$;*
(2) *$[b,T]$ is a bounded operator from $M^p_q$ to $WM^s_t$.*

**Theorem 3.7.** *Let $0<\alpha<1$, $1<q\leq p<\infty$, $1<t\leq s<\infty$, $1<l\leq s<\infty$, $\frac{1}{s}=\frac{1}{p}-\frac{\alpha}{n}$, $\frac{1}{t}=\frac{1}{q}-\frac{\alpha}{n}$ and $\frac{l}{s}=\frac{q}{p}$,. The following statements are equivalent:*

(1) *$b\in \mathrm{Lip}_\alpha$;*
(2) *$[b,T]$ is a bounded operator from $M^p_q$ to $M^s_l$;*
(3) *$[b,T]$ is a bounded operator from $M^p_q$ to $M^s_t$;*
(4) *$[b,T]$ is a bounded operator from $M^p_q$ to $WM^s_t$.*

*Proof of Theorem 3.6.* (1) $\Rightarrow$ (2): For $q > 1$, Theorem 3.4 and (3.1) yield

$$\|[b,T]f\|_{WM^s_t} \le \|b\|_{\mathrm{Lip}_\alpha}\|I_\alpha(|f|)\|_{WM^s_t} \le \|b\|_{\mathrm{Lip}_\alpha}\|I_\alpha(|f|)\|_{M^s_t} \le C\|b\|_{\mathrm{Lip}_\alpha}\|f\|_{M^p_q}.$$

For $q = 1$, it follows from [11, Theorem 1.4] that

$$\|I_\alpha(f)\|_{WM^s_1} \le C\|f\|_{M^p_q},$$

which gives

$$\|[b,T]f\|_{WM^s_1} \le \|b\|_{\mathrm{Lip}_\alpha}\|I_\alpha(|f|)\|_{WM^s_1} \le C\|b\|_{Lip_\alpha}\|f\|_{M^p_q}.$$

(2) $\Rightarrow$ (1): A same argument as the proof of Theorem 2.2 except choosing $q_1 \in (1,t)$ and replacing (2.1) by

$$\begin{aligned}
\frac{1}{|Q|}\int_Q |b(x) - b_Q|dx &\le \frac{2}{|Q|}\int_Q |b(x) - b_{Q'}|dx \\
&\le \frac{C}{|Q|^{1/s}}\sum |a_m|\|[b,T](g_m)\|_{M^s_{q_1}} \\
&\le \frac{C}{|Q|^{1/s}}\sum |a_m|\|[b,T](g_m)\|_{WM^s_t} \\
&\le C|Q|^{\alpha/n}\|[b,T]\|_{M^p_q \to WM^s_t}.
\end{aligned}$$

implies $b \in \mathrm{Lip}_\alpha$ due to Lemma 3.1. □

*Proof of Theorem 3.7.* (1) $\Rightarrow$ (2): It follows from (3.1) and Theorem 3.5 that $[b,T]$ is bounded from $M^p_q$ to $M^s_l$.

(2) $\Rightarrow$ (3): Since $\frac{1}{s} = \frac{1}{p} - \frac{\alpha}{n}$, $\frac{1}{t} = \frac{1}{q} - \frac{\alpha}{n}$ and $\frac{l}{s} = \frac{q}{p}$, we have

$$\frac{1}{q} - \frac{1}{t} = \frac{1}{p} - \frac{1}{s} = \frac{q}{p}\Big(\frac{1}{q} - \frac{1}{l}\Big) \le \frac{1}{q} - \frac{1}{l},$$

and hence $t \le l$. Then $\|\cdot\|_{M^s_t} \le \|\cdot\|_{M^s_l}$.

It is obvious for (3) $\Rightarrow$ (4).

(4) $\Rightarrow$ (1): By Theorem 3.6, we obtain $b \in \mathrm{Lip}_\alpha$. Hence, the proof of Theorem 3.7 is completed. □

## References


[1] D. R. Adams, *A note on Riesz potentials*, Duke Math. J. **42** (1975), 765-778.

[2] R. Coifman, R. Rochberg and G. Weiss, *Factorization theorems for Hardy spaces in several variables*, Ann. of Math. **103** (1976), 611-635.

[3] R. A. Devore and R. C. Sharpley, *Maximal functions measuring smoothness*, Mem. Amer. Math. Soc. **47** (1984).

[4] Y. Ding, *A characterization of BMO via commutators for some operators*, Northeast. Math. J. **13** (1997), 422-432.

[5] S. Janson, *Mean oscillation and commutators of singular integral operators*, Ark. Math. **16** (1978), 263-270.

[6] S. Janson, M. Taibleson and G. Weiss, *Elementary characterization of the Morrey-Campanato spaces*, Lect. Notes in Math. **992** (1983), 101-114.

[7] H. Kozono and M. Yamazaki, *Semilinear heat equations and the Navier-Stokes equation with distributions in new function spaces as initial data*, Comm. Partial Differential Equations **19** (1994), 959-1014.

[8] M. Paluszyński, *Characterization of the Besov spaces via the commutator operator of Coifman, Rochberg and Weiss*, Indiana Univ. Math. J. **44** (1995), 1-17.

[9] J. Peetre, *On the theory $\mathcal{L}_{p,\lambda}$ spaces*. J. Funct. Anal. **4** (1969), 71-87.

[10] S. G. Shi and S. Z. Lu, *Some characterizations of Campanato spaces via commutators on Morrey spaces*, Pacific J. Math. **264** (2013), 221-234.

[11] L. Tang, *Endpoint estimates for multilinear fractional integrals*, J. Aust. Math. Soc. **84** (2008), 419-429.

College of Mathematics and System Sciences
Xinjiang University
Urumqi 830046
Republic of China
*E-mail address*: Wangdh1990@126.com; zhoujiangshuxue@126.com

School of Mathematics and Statistics
WuHan University
WuHan 430072
Republic of China
*E-mail address*: wychencn@hotmail.com